ANTERIOR AND POSTERIOR LUMBAR

SURGERY Name Date:

PREADMISSION INSTRUCTION SHEET

PLEASE READ CAREFULLY

You are scheduled for surgery at Community Hospital, (date) (day) , (arrival time)

The hospital is prepared for you to have your pre-op lab work, chest x-ray and ekg done within seven (7) days of the procedure. This can be done on a walk in basis at the outpatient entrance to the facility, which can be found at the back of the building by following the signs past the emergency room.

Please notify the nurse if you are taking a blood thinner such as asprin, Plavix, or Coumadin. You should stop taking most blood thinners up to five days before your procedure and should discuss this with the nurse. It is your responsibility to check with the doctor that has put you on this medication to ensure you are medically stable to be off from it for the required time prior to surgery.

The hospital will instruct you on taking other medication prior to surgery.

Due to the type of surgery you will be undergoing Dr Alvis has very specific instructions in relation to your diet starting at noon the day before surgery. Failure to follow them carefully may force cancellation of your surgery.

DO NOT EAT ANYTHING AFTER 12
NOON THE DAY BEFORE SURGERY.
YOU MAY ONLY HAVE CLEAR LIQUIDS
SUCH AS COFFEE, TEA, COLA, JELLO,
BROTH.
6:00 PM DRINK ONE 8 OUNCE BOTTLE OF
MAGNESIUM CITRATE. THIS CAN BE
FOUND AT ANY PHARMACY.
12PM (MIDNIGHT) THE NIGHT BEFORE
SURGERY DO NOT EAT ANYTHING ELSE
BY MOUTH.

COMMUNITY HOSPITAL 3100 SW 89.111 STREET ON 89^{Tit} BETWEEN MAY AND 144 (405) 602-8100

DR ALVIS2412 Palmer Circle **NORMAN, OK 73069**(405) 321-6347

WOUND CARE

24 hours after surgery (if you are home from the hospital) you may remove the dressings to your lower back and/or abdomen. The incisions should be left open to air as much as possible.

The incisions may have areas of drainage or bleeding. This is normal. Please cover the area that is draining with a small dressing to protect your clothing. DO NOT COVER THE ENTIRE INCISION WITH A DRESSING. Change the dressing every day if necessary.

The incisions need to be washed daily with soap and water once the dressing is removed. This does not require special soap. The best way to do this is in the shower. Be sure to avoid the shower spray hitting the incisions directly.

DO NOT TAKE A TUB BATH UNTIL AFTER YOUR TWO WEEK FOLLOW UP APPOINTMENT. DO NOT SUBMURGE THE INCISION

You may apply ice to the incision as needed to aid with swelling.

RESTRICTIONS

It is not recommended that you drive for 10-14 days after surgery depending on your pain level and the amount of medication needed for pain control.

Walk as tolerated. You are not bed bound or home bound *as a* result of this surgery. You are encouraged to increase your activity as tolerated. Rest when your body indicates you need to rest. You may find that you have a good day and excel that day, and then the next day you do not feel *as* well. This is normal and to be expected as you heal.

MEDICATION

Take your pain medication as prescribed. The first two weeks you may have increased pain due to the swelling from surgery and you may need to take your medication on a schedule.

DO NOT TAKE YOUR MEDICATION IF IT IS NOT DUE. If your medication is not working YOU are responsible for contacting the prescription line at the office to request a change. You may have an increase in pain after surgery, which is to be expected, but we can make changes to help with the pain if necessary. Keep your medications secure. It is your responsibility to protect your medications. Typically, if your coeds are lost, stolen, or used up too soon they will not be replaced. If you have any question about this, please contact the prescription line.

Dr Alvis is not in the office every day and therefore we need you to give us several days notice if you are going to run out of your medication. Many of the medications that he prescribes can only be dispensed on a written prescription and will have to be picked up at the office. We need your assistance in making sure you do not run out of medication.